

SECTION 1

INTRODUCTION

1.1 This manual was prepared to assist biologists and managers in USEPA and other Federal, state, and private water monitoring organizations in the use of fish as indicators of ecosystem health and for evaluating the biological integrity of surface waters and protecting quality water resources. The manual contains biological criteria and laboratory and field methods that will aid in the monitoring and bioassessment of the effects of anthropogenic and environmental stresses on fish populations and communities. It will also facilitate the expansion and refinement of our knowledge of the ecological requirements of fish species in freshwater, estuarine, and marine habitats.

1.2 The manual includes sections on quality assurance and quality control, safety and health, sampling methods and techniques, sample preservation and identification, data analyses, special techniques, bioassessment protocols for use in streams and rivers, a family-level ichthyoplankton index method, fish health and condition assessment procedures, guidelines for fish sampling and tissue preparation for bioaccumulative contaminants, and a fisheries bibliography. Guidelines and procedures for fish kill investigations are provided.

1.3 Fish community evaluation and assessment should measure the overall structure (number of species and individuals within a community) and function (organism interaction in the utilization of food and other biological resources) of various aquatic habitats considered for study. These measurements should include such factors as habitat characteristics and quality, riparian vegetation, and hydraulic characteristics that are expected to influence fish community spatial and temporal variability. One must also distinguish the alterations induced by anthropogenic activities from natural variations which occur in the environment.

1.4 In North America, fish are the focus of economically important sport and commercial fisheries, and are an important source of food for humans. To the general public the size and species composition of a fish community is the most meaningful index of pollution.

1.5 In most aquatic ecosystems, fish are usually the most common vertebrates. Fish communities occupy the upper trophic levels of aquatic food webs, and they are dependent on the same or other trophic level life forms for food. In aquatic communities fish can be one of the most sensitive indicators of water quality assessment and biological integrity in aquatic environments (Angermeier et al., 1991; Fausch et al., 1990; Karr, 1981, 1987, 1990, 1991; Smith, 1971; McKenzie et al., 1992). The literature contains much data on fish species distribution, life histories, ecology, pollution tolerance, and environmental requirements. Fish are directly and indirectly affected by chemical and physical changes in the environment, and the population or community of fish in rivers, streams, lakes, estuaries, and oceans reflects the state of the health of the aquatic environment or watershed as a whole.

Because they are conspicuous, fish populations or fish assemblages are commonly used as environmental indicators or as an index for water quality (Table 1).

1.6 Water quality conditions that significantly affect the lower levels of food webs (e.g., plankton and benthic invertebrates, including macroinvertebrates, USEPA, 1990a) will affect the abundance and species composition of the fish population. In some cases, fish may exhibit signs of being more sensitive to certain pollutants than are the lower animals and plants, and may be adversely affected even when the lower levels of food webs are relatively unharmed.

1.7 Karr (1981, 1987), Karr et al. (1986, 1987), Ohio EPA (1990), and USEPA (1990a,b) have indicated that five major sets of abiotic and biotic factors affect and ascertain biological integrity or water resource integrity (Table 2). To determine anthropogenic or natural impact on aquatic ecosystems, all monitoring or bioassessment programs must survey and evaluate in a methodical and systematic way all five sets of factors. Although a thorough discussion of all these factors is beyond the scope of this document, a discussion of how some of these factors influence the biological integrity of surface waters and several methods and procedures in evaluating these complex set of factors are presented here. For a more comprehensive discussion of all these factors, consult USEPA (1990a, 1990b), Ohio EPA (1990), and the references in Section 12, Fisheries Bibliography.

1.8 Many species of fish have stringent dissolved oxygen and temperature requirements and are intolerant to chemical and physical contaminants resulting from municipal, agricultural, industrial, forestry, and mining activities. Also, fish communities are sensitive to and good indicators of macrohabitat disturbances (Rankin, 1989).

1.9 The discharge of moderate amounts of degradable organic wastes may increase the nutrient levels (eutrophication) in the habitat and result in an increase in the standing crop (total amount of the biomass of organisms of one or more species within a locality) of fish. This increase usually occurs in one or a few species and results in an imbalance in the population. The discharge of large amounts of degradable organic materials may result in depressed oxygen levels which may reduce the number and kinds of fishes present and increase the standing crop of pollution tolerant species. In extreme cases the fishery may be eliminated in the affected area.

1.10 The effects of toxic wastes may range from the elimination of most fish to a reduction in reproductive capacity (fecundity) or resistance to disease and parasitism. Massive and complete fish kills are dramatic signs of abrupt, adverse changes in environmental conditions. Fish, however, can repopulate an area rapidly if the habitat is not destroyed and the water quality improves. The cause of the fish kill may be difficult to detect by examination of the fish community after it has recovered from the effects of the pollutant. Chronic pollution, on the other hand, is more selective in its effects, exerts its influence over a long period of time, and causes recognizable changes in the species composition and relative abundance of the fish.

TABLE 1. ATTRIBUTES OF FISHES AND DESIRABLE COMPONENTS FOR BIOASSESSMENT AND BIOMONITORING PROGRAMS¹

Goal/Quality	Attribute
Accurate Assessment of Aquatic Ecosystem Integrity	Fish populations and individuals generally remain in the same area during summer seasons.
	Communities are persistent and usually recover rapidly from natural disturbances. Comparable results can be expected from an unperturbed site at various times within a season.
	Fish have larger home ranges and are less affected by natural microhabitat differences than smaller organisms, such as macroinvertebrates. This makes fish extremely useful for assessing regional, macrohabitat, and mesohabitat differences.
	Most fish species have long life spans (3-10+ years) and can reflect both long term and current water resource quality.
	Fish continually inhabit the receiving water and reflect the chemical, physical, and biological histories of the water.
Visibility	Fish represent a broad spectrum of community tolerances from very sensitive to highly tolerant, and respond to chemical, physical, and biological degradation in characteristics response patterns.
	Fish are a highly visible component of the aquatic community, and so are of interest to the public.
Ease of Use and Interpretation	Aquatic resource uses and regulatory language are generally characterized in terms of fish (i.e., fishable and swimmable goals of the Clean Water Act).
	The sampling frequency for trend assessment is less than for short-lived organisms.
	The taxonomy of fishes is well established, allowing professional biologists the ability to reduce laboratory time by identifying many specimens in the field.
	The distribution, life histories, and tolerances to environmental stresses of most North American species are well documented in the literature.

¹Adapted from Simon (1991).

TABLE 2. FIVE MAJOR CLASSES OF ENVIRONMENTAL FACTORS WHICH INFLUENCE AND DETERMINE THE BIOLOGICAL INTEGRITY OF SURFACE WATERS WITH SOME OF THEIR IMPORTANT CHEMICAL, PHYSICAL, AND BIOLOGICAL COMPONENTS IN LENTIC AND LOTIC SYSTEMS¹

1. ENERGY SOURCE

STREAMS, RIVERS

Nutrient cycling
Organic matter particle size
Primary productivity
Seasonal cycles
Solar radiation

LAKES, RESERVOIRS, ESTUARIES, OCEANS

Nutrients cycling
Organic matter particle size
Primary productivity
Seasonal cycles
Solar radiation

2. WATER QUALITY/CHEMICAL VARIABLES

STREAMS, RIVERS

Adsorption
Alkalinity
DO
Hardness
Metals, other toxic substances
Nutrients
Organics
pH
Solubility
Temperature
Turbidity
Water cycling

LAKES, RESERVOIRS, ESTUARIES, OCEANS

Adsorption
Alkalinity
DO
Hardness
Metals, other toxic substances
Nutrients
Organics
pH
Solubility
Temperature
Turbidity
Water cycling

3. HABITAT QUALITY

STREAMS, RIVERS

Bank stability
Canopy
Channel morphology (riffles, pools)
Current velocity
Gradient
Instream cover (woody debris)
Riparian vegetation
Siltation
Sinuosity
Substrate types
Width/depth

LAKES, RESERVOIRS, ESTUARIES, OCEANS

Bank stability
Shoreline vegetation
Substrate types
Siltation
Wave action
Width/depth
Inwater abiotic/biotic cover

¹Adapted from Karr (1987, 1991), Karr and Dudley (1981), Karr et al. (1986, 1987), and USEPA (1990a; 1990b).

TABLE 2. FIVE MAJOR CLASSES OF ENVIRONMENTAL FACTORS WHICH INFLUENCE AND DETERMINE THE BIOLOGICAL INTEGRITY OF SURFACE WATERS WITH SOME OF THEIR IMPORTANT CHEMICAL, PHYSICAL, AND BIOLOGICAL COMPONENTS IN LENTIC AND LOTIC SYSTEMS (CONTINUED)

4. FLOW REGIME

STREAMS, RIVERS

Ground water
High/low extremes
Land use
Precipitation/runoff
Water volume

LAKES, RESERVOIRS, ESTUARIES, OCEANS

Ground water
High/low extremes
Land use
Precipitation/runoff
Water volume

5. BIOTIC ASSOCIATIONS

STREAMS, RIVERS

Feeding
Competition
Disease
Parasitism
Predation
Reproduction

LAKES, RESERVOIRS, ESTUARIES, OCEANS

Feeding
Competition
Disease
Parasitism
Predation
Reproduction

1.11 The utilization of biological components (structural and functional) to evaluate the ambient aquatic community of our nations surface water has been discussed and well documented in the literature. Some recent examples are Crowder (1990), Downing et al. (1990), Fausch et al. (1990), Hunsaker and Carpenter (1990), Karr et al. (1986), Karr, (1991), Ohio EPA (1987a, 1987b, 1989, 1990), Plafkin et al. (1989), Shuter (1990), Simon (1991), and USEPA (1990a, 1990b). Structural components of fish communities include diversity, taxa guilds, numbers, and biomass. Functional components of fish communities include the feeding or trophic strategy, reproductive behavior and guild classification, and environmental tolerance to perturbations.

1.12 The principal characteristics of interest in bioassessment studies of fish populations include: (1) species richness (number of species)--presence or absence; relative and absolute abundance of each species, (2) size distribution, (3) habitat guilds--pelagic, littoral, and benthic species, (4) trophic guilds--omnivores, piscivores, and invertivores, (5) growth rate, (6) condition factor, (7) reproductive guilds, egg production and success, (8) general tolerance guilds (indicator taxa)--intolerant, tolerant, and sensitive species, (9) incidence of disease and parasitism (10) fish kills, (11) palatability, and (11) fishability--catchability, desirability, and sustainability. Observations of fish behavior can also be valuable in detecting environmental problems, e.g., ventilation rates, position in the current, and erratic movement. Fish may also be utilized for field and laboratory bioassays (USEPA, 1991a, 1991b, 1992a, 1992b), for tissue analyses to measure the concentrations of metals and pesticides (see Section 10, Guidelines for Fish Sampling and Tissue Preparation for bioaccumulative Contaminants) for histopathologic examination (Hinton and Lauren, 1990), and biomarker studies (Adams, 1990a, 1990b; Anderson, 1990; Jimenez and Stegeman, 1990; Rice, 1990; Schreck, 1990; and Thomas, 1990).

1.13 Fisheries data are useful in enforcement cases and in long-term water quality status and trends monitoring (Tebo, 1965; Ohio EPA, 1990; USEPA, 1991a). Before fishery surveys are initiated, a careful and exhaustive search should be conducted for existing information on the fish populations or communities in question. State and Federal fishery agencies and universities may be potential sources of information. If data are not available and a field study must be conducted, State and other Federal agencies may assist in a survey and may provide needed expertise and specialized equipment for the collection of specific, local fishes. A joint effort is usually more economical and efficient and will promote continued cooperation between agencies and parties involved.

1.14 Fisheries data may have limitations. Even if the species composition of the fish in a specific area is known before and after the discharge of pollutants, the significance of changes in the catch might not be satisfactorily interpreted unless there are adequate data on spawning, seasonal migration, temperature requirements and stream-flow responses, feeding activities, diurnal movements, habitat preferences, and activity patterns. Without adequate data, fish presence or absence cannot be directly correlated with water quality. Furthermore, any existing data of known quality on the water quality requirements of fish would be of value in interpreting field data.

1.15 Federal and state regulations usually require a fish collecting permit because some species of fish are protected by law, and the collection of others is regulated. The state fishery agencies must be contacted before fish can be taken in a field study. Investigators should confirm that they have complied with federal and state regulations before collecting samples of fish. The state should be contacted prior to any fish study to ensure that investigators comply with current regulations.

1.16 The design of fish studies should be based upon study goals and data quality objectives (DQOs) (see Section 2, Quality Assurance and Quality Control). To supplement the material contained in this manual, a number of basic references should be reviewed by investigators involved in fish sampling programs and studies. Useful references include Adams (1990), Angermeier et al. (1991), APHA (1992), Bartell (1990), Edwards and Megrey (1989), Evans et al. (1990), Everhart and Youngs (1981), Fausch et al. (1990), Gammon (1980), Gammon et al. (1990), Hankin and Reeves (1988), Hellawell (1986), Herricks and Schaeffer (1985), Hirsch et al. (1988), Hughes et al. (1986), Johnson and Nielsen (1983), Karr (1981, 1987, 1990, 1991), Karr and Dionne, 1991, Karr and Dudley (1981), Karr et al. (1983, 1986, 1987), Magnuson (1991), Mancini (1989), Mangel and Smith (1990), Minshall et al. (1989), Ohio EPA (1986, 1987a, 1987b, 1989, 1990), Omernik (1987), Platts et al. (1983), Robins et al. (1991), Schreck and Moyle (1990), Templeton (1984), Tonn (1990), USEPA (1988), USEPA (1990a, 1990b), (USEPA, 1991c, 1991d, 1991e), Whittier and Paulsen (1992), Wooten (1990), and Yoder (1991).

1.16.1 If fish data are to be useful, they must be acquired according to standardized sampling methods and analyzed with appropriate statistical methods. Two very important qualities of sampling data are accuracy and precision. Accuracy refers to how well the sample represents the whole of the study. In fishery studies, collecting accurate (or unbiased) data may be difficult because studies are poorly designed. Precision refers to repeatability of data. To supplement the statistics in this document, investigators should consult the commonly cited statistical references (Cochran, 1977; Conover, 1980; Green, 1979; Hicks, 1982; Snedecor and Cochran, 1981; Sokal and Rohlf, 1981; Zar, 1984).

1.17 Literature Cited

- Adams, S.M. (ed.). 1990a. Biological indicators of stress in fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD.
- Adams, S.M. 1990b. Status and use of biological indicators for evaluating the effects of stress on fish. In: Adams, S.M. (ed.). Biological indicators of stress in fish. American Fisheries Society, Symposium 8, American Fisheries Society, Bethesda, MD. pp. 1-8.
- Anderson, D.P. 1990. Immunological indicators: Effects of environmental stress on immune protection and disease outbreak. In: Adams, S.M. Biological indicators of stress in fish. American Fisheries Society, Symposium 8, American Fisheries Society, Bethesda, MD. pp. 38-50.

- Angermeier, P.L., R.J. Neves, and L.A. Nielsen. 1991. Assessing stream values: Perspectives of aquatic resource professionals. *North Amer. J. Fisheries Management* 11(1):1-10.
- APHA. 1992. Standard methods for the examination of water and wastewater. American Public Health Association, American Water Works Association, and Water Pollution Control Federation, (18th ed.), Washington, DC.
- Bartell, S.M. 1990. Ecosystem context for estimating stress-induced reductions in fish populations. *In*: Adams, S.M. (ed.). Biological indicators of stress fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD. pp. 167-182.
- Cochran, W.G. 1977. Sampling techniques. John Wiley and Sons, Inc., New York, NY.
- Conover, W.J. 1980. Practical nonparametric statistics. John Wiley, New York, NY.
- Crowder, L.B. 1990. Community ecology. *In*: Schreck, C.B. and P.B. Moyle (eds.). Methods for fish biology. Amer. Fish. Soc., Bethesda, MD. pp. 609-632.
- Downing, J.A., C. Plante, and S. Lalonde. 1990. Fish production correlated with primary productivity, not the morphoedaphic index. *Can. J. Fish. Aquatic Sci.* 47(10):1929-1936.
- Edwards, E.F. and B.A. Megrey (eds.). 1989. Mathematical analysis of fish stock dynamics. American Fisheries Symposium 6, American Fisheries Society, Bethesda, MD.
- Evans, D.O., G.J. Warren, V.W. Cairns. 1990. Assessment and management of the fish community health in the Great Lakes: Synthesis and recommendation. *J. Great Lakes Res.* 16(4):639-669.
- Everhart, W.H. and W.D. Youngs. 1981. Principles of fishery science. Cornell University Press, Ithaca, NY.
- Fausch, K.D., J. Lyons, J.R. Karr, and P.L. Angermeier. 1990. Fish communities as indicators of environmental degradation. *In*: Adams, S.M. (ed.). Biological indicators of stress fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD. pp. 123-144.
- Gammon, J.R. 1980. The use of community parameters derived from electrofishing catches of river fish as indicators of environmental quality. *In*: Seminar on Water Quality Management Tradeoffs. EPA-905/9-80-009, U.S. Environmental Protection Agency, Washington, DC.
- Gammon, J.R., C.W. Gammon, and M.K. Schmid. 1990. Land use influence on fish communities in central Indiana streams. *In*: W.S. Davis (ed.). Proceedings of the 1990 Midwest Pollution Control biologists Meeting.

EPA 905/9-90-995, U.S. Environmental Protection Agency, Chicago, IL.
pp. 111-120.

- Green, R.H. 1979. Sampling design and statistical methods for environmental biologists. John Wiley, New York, NY.
- Hankin, D.G. and G. H. Reeves. 1988. Estimating total fish abundance and total habitat area in small streams based on visual estimation methods. *Can. J. Fish. Aquat. Sci.* 45(5):834-844.
- Hellawell, J.M. 1986. Biological indicators of freshwater pollution and environmental management. Elsevier Science Publishing, Co., Inc., New York, NY
- Herricks, E.E. and D.J. Schaeffer. 1985. Can we optimize biomonitoring? *Env. Mgmt.* 9:487-492.
- Hicks, C.R. 1982. Fundamental concepts in the design of experiments. Holt, Rinehart, and Winston, New York, NY.
- Hinton, D.E. and D.J. Lauren. 1990. Integrative histopathological approaches to detecting effects of environmental stressors on fishes. *In: Adams, S.M. (ed.). Biological indicators of stress in fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD.* pp. 51-66.
- Hirsch, R.M., W.M. Alley, and W.G. Wilber. 1988. Concepts for a national water-quality assessment program. U.S. Geological Survey Circular 1021, Federal center, Denver, CO.
- Hughes, R.M., D.P. Larsen, and J.M. Omernik. 1986. Regional reference sites: a method for assessing stream pollution. *Env. Mgmt.* 10(5):629-635.
- Jimenez, B.D. and J.J. Stegeman. 1990. Detoxication enzymes as indicators of environmental stress on fish. *In: Adams, S.M. (ed.). Biological indicators of stress fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD.* pp. 67-79.
- Johnson, D.L. and L.A. Nielsen. 1983. Sampling considerations. *In: Nielsen, L.A. and D.L. Johnson (eds.). Fisheries Techniques. American Fisheries Society, Bethesda, MD.* pp. 1-21.
- Karr, J.R. 1981. Assessment of biotic integrity using fish communities. *Fisheries* 6(6):21-27.
- Karr, J.R. 1987. Biological monitoring and environmental assessment: a conceptual framework. *Environmental Management* 11:249-256.
- Karr, J.R. 1990a. Biological integrity and the good of environmental legislation: Lessons for conservation biology. *Conservation Biology* 4:244-250.

- Karr, J.R. 1990b. Bioassessment and non-point source pollution: an overview. Pages 4-1 to 4-18. *In*: Second National Symposium on Water Quality Assessment. Meeting summary, October 16-19, 1989, Fort Collins, Colorado, U.S. Environmental Protection Agency, Washington, DC.
- Karr, J.R. 1991. Biological integrity: A long-neglected aspect of water resource management. *Ecological Applications* 1:66-84.
- Karr, J.R. and M. Dionne. 1991. Designing surveys to assess biological integrity in lakes and reservoirs. *In*: Biological Criteria: Research and Regulation. Proceedings of a symposium, pp. 62-72, EPA/440/5-91-005. U.S. Environmental Protection Agency, Office of Water, Washington, DC.
- Karr, J.R. and D.R. Dudley. 1981. Ecological perspective on water quality goals. *Env. Mgmt.* 5:55-68.
- Karr, J.R., K.D. Fausch, P.L. Angermeier, P.R. Yant, and I.J. Schlosser. 1986. Assessing biological integrity in running waters: a method and its rationale. Special Publication 5. Illinois Natural History Survey, Urbana, IL.
- Karr, J.R., L.A. Toth, and G.D. Garman. 1983. Habitat preservation for midwest stream fishes: principles and guidelines. EPA-600/3-83-006. U.S. Environmental Protection Agency, Corvallis, OR.
- Karr, J.R., P.R. Yant, K.D. Fausch, and I.J. Schlosser. 1987. Spatial and temporal variability of the index of biotic integrity in three midwestern streams. *Trans. Amer. Fish. Soc.* 116:1-11.
- Magnuson, J.J. 1991. Fish and fisheries ecology. *Ecol. Application* 1(1):13-26.
- Manci, K.M. 1989. Riparian ecosystem creation and restoration: A literature summary. Fish and Wildlife Service, U.S. Dept. Interior, Washington, DC.
- Mangel, M. and P.E. Smith. 1990. Presence-absence sampling for fisheries management. *Can. J. Fish. Aquat. Sci.* 47:1875-1887.
- McKenzie, D.H., D.E. Hyatt, and V.J. McDonald (eds.). 1992. Ecological Indicators. Proceedings of an International Symposium, Fort Lauderdale, USA, October 16-19, 1990. Volume I and II. Elsevier Applied Science, Elsevier Publishers, Ltd., London and New York.
- Minshall, G.W., S.E. Jensen, and W.S. Platts. 1989. The ecology of stream and riparian habitats of the Great Basin region: A community profile. National Wetlands Research Center, U.S. Fish and Wildlife Service, Slidell, LA.

- Ohio EPA. 1986. The cost of biological field monitoring. Ohio Environmental Protection Agency, Division of Water Qual. Monitoring and Assessment, Evaluation and Standards Section, Columbus, OH.
- Ohio EPA. 1987a. Biological Criteria for the protection of aquatic life: Volume I: The role of biological data in water quality assessment. Ohio Environmental Protection Agency, Division of Water Quality Planning and Assessment, Ecological Assessment Section, Columbus, OH.
- Ohio EPA. 1987b. Biological criteria for the protection of aquatic life: Volume II. Users manual for biological field assessment of Ohio surface waters. Ohio Environmental Protection Agency, Division of Water Quality Planning and Assessment, Ecological Assessment Section, Columbus, OH.
- Ohio EPA. 1989. Biological criteria for the protection of aquatic life: Volume III. Standardized biological field sampling and laboratory methods for assessing fish and macroinvertebrate communities. Ohio Environmental Protection Agency, Division of Water Quality Planning and Assessment, Ecological Assessment Section, Columbus, OH.
- Ohio EPA. 1990. The use of biocriteria in the Ohio EPA surface water monitoring and Assessment Program. Ohio Environmental Protection Agency, Division of Water Quality Planning and Assessment, Ecological Assessment Section, Columbus, OH.
- Omernik, J.M. 1987. Ecoregions of the conterminous United States. *Ann. Assoc. Amer. Geogr.* 77:117-125.
- Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross, and R.M. Hughes. 1989. Rapid bioassessment protocols for use in streams and rivers: benthic macroinvertebrates and fish. EPA/440/4-89/001. U.S. Environmental Protection Agency, Assessment and Watershed Protection Division, Washington, DC.
- Platts, W.S., W.F. Megahan, and G.W. Minshall. 1983. Methods for evaluating streams, riparian, and biotic conditions. General Technical Report INT-138, Intermountain Forest and Range Experiment Station, Forest Service, U.S. Dept. Agriculture, Ogden, UT.
- Rankin, E.T. 1989. The qualitative habitat evaluation index (QHEI): rationale, methods, and application. Ohio Environmental Protection Agency, Division Water Quality, Planning and Assessment, Ecological Assessment Section, P.O. Box 1049, 1800 WaterMark Drive, Columbus, OH.
- Rice, J.A. 1990. Bioenergetics modeling approaches to evaluation of stress in fish. In: Adams, S.M. (ed.). Biological indicators of stress fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD. pp. 80-92.
- Robin, C.R., C.E. Bond, J.R. Brooker, E.A. Lachner, R.N. Lea, and W.B. Scott. 1991. Common and scientific names of fishes from the United States and

- Canada. Fifth Edition. American Fisheries Society, Special Publication 20, American Fisheries Society, Bethesda, MD.
- Schreck, C.B. 1990. Physiological, behavioral, and performance indicators of stress. *In*: Adams S.M. (ed.). Biological indicators of stress fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD. pp. 29-37.
- Schreck, C.B. and P.B. Moyle (eds.). 1990. Methods for fish biology. American Fisheries Society, Bethesda, MD.
- Shuter, B.J. 1990. Population-level indicators of stress. *In*: Adams, S.M. (ed.). Biological indicators of stress in fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD. pp. 145-166.
- Simon, T.P. 1991. Development of index of biotic integrity expectations for the ecoregions of Indiana. I. Central Corn Belt Plain. EPA-905/9-91/025. U.S. Environmental Protection Agency, Environmental Science Division, Monitoring and Quality Assurance Branch, Ambient Monitoring Section, Chicago, IL.
- Smith, P.W. 1971. Illinois streams: a classification based on their fishes and an analysis of factors responsible for the disappearance of native species. Ill. Nat. Hist. Surv. Notes 76.
- Snedecor, G.W. and W.G. Cochran. 1981. Statistical methods, Iowa State University Press, Ames, IA.
- Sokal, R.R. and F.J. Rohlf. 1981. Biometry, Freeman, San Francisco, CA.
- Tebo, Jr., L.B. 1965. Fish population sampling studies at water pollution surveillance system stations on the Ohio, Tennessee, Clinch, and Cumberland Rivers. Applications and development Report No. 15, Div. Water Supply and Pollution Control, U.S. Public Health Service, Cincinnati, OH.
- Templeton, R.G. 1984. Freshwater fisheries management. Fishing News Books, Ltd., Farnham, Surrey, England, U.K.
- Thomas, P. 1990. Molecular and biochemical responses of fish to stressors and their potential use in environmental monitoring. *In*: Adams, S.M. (ed.). Biological indicators of stress fish. American Fisheries Symposium 8, American Fisheries Society, Bethesda, MD. pp. 9-28.
- Tonn, W.M. 1990. Climate change and fish communities: A conceptual framework. Trans. Amer. Fish. Soc. 119:337-352.
- USEPA. 1988. The lake and reservoir restoration guidance manual. EPA 440/5-88-002. U.S. Environmental Protection Agency, Criteria and Standards Division, Nonpoint Sources Branch, Washington, DC.

- USEPA. 1990a. Macroinvertebrate field and laboratory methods for evaluating the biological integrity of surface waters. Donald J. Klemm, Philip A. Lewis, Florence Fulk, and James M. Lazorchak. EPA/600/4-90/030. U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory, Cincinnati, OH.
- USEPA. 1990b. Biological Criteria. National Program guidance for Surface Waters. EPA/440/5-90/004. U.S. Environmental Protection Agency, Office of Water, Criteria and Standards Division, Office of Water Regulations and Standards, Washington, DC.
- USEPA. 1991a. Technical support document for water quality-based toxics control. EPA/5052-90/001. U.S. Environmental Protection Agency, Office of Water Enforcement and Permits and Office of Water Regulations and Standards, Washington, DC.
- USEPA. 1991b. Methods for measuring the acute toxicity of effluents and receiving waters to freshwater and marine organisms. Cornelius I. Weber (ed.). Fourth Edition. EPA/600/4-90/027. U.S. Environmental Protection Agency, Monitoring Systems Laboratory, Cincinnati, OH.
- USEPA. 1991c. Biological Criteria. State development and implementation efforts. EPA-440/5-91-003. U.S. Environmental Protection Agency, Office of Water, Washington, DC.
- USEPA. 1991d. Biological criteria. Guide to technical literature. EPA-440/5-91-004. U.S. Environmental Protection Agency, Office of Water, Washington, DC.
- USEPA. 1991e. Biological criteria: Research and regulation. EPA-440/5-91-005. U.S. Environmental Protection Agency, Office of Water, Washington, DC.
- USEPA. 1992a. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to marine and estuarine organisms. Donald J. Klemm and George E. Morrison (eds.). Second Edition. EPA/600/4-91-021. U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory, Cincinnati, OH.
- USEPA. 1992b. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms. Philip A. Lewis, Donald J. Klemm, and James M. Lazorchak (eds.). Third Edition. EPA/600/4-91/022. U.S. Environmental Protection Agency, Environmental Monitoring Systems Laboratory, Cincinnati, OH.
- Whittier, T.R. and S.G. Paulsen 1992. The surface waters component of the Environmental Monitoring and Assessment Program (EMAP): an overview. J. Aquatic Ecosystem Health 1:119-126.
- Wooten, R.J. 1990. The ecology of teleost fishes. Chapman and Hall Press, New York, NY.

- Yoder, C.O. 1991. The integrated biosurvey as a tool for evaluation of aquatic life use attainment and impairment in Ohio surface waters. *In: Biological Criteria: Research and Regulation. Proceedings of a Symposium.* EPA/440/5-91-005. U.S. Environmental Protection Agency, Office of Water, Washington, DC. pp. 110-122.
- Zar, J. H. 1984. Biostatistical analysis. Prentice-Hall, Inc., Englewood Cliffs, NJ.